

Fermentation of seaweed flour with various fermentors to improve the quality of fish feed ingredients

Fermentasi tepung rumput laut dengan berbagai fermentor untuk meningkatkan kualitas sebagai bahan baku pakan ikan

Siti Aslamyah*, Muh. Yusri Karim, Badraeni

Program Studi Budidaya Perairan, Fakultas Ilmu Kelautan dan Perikanan
Universitas Hasanuddin, Jalan Perintis Kemerdekaan Km X, Tamalanrea, Makassar 90245
*E-mail: siti_aslamyah_uh@yahoo.co.id/sitiaslamyah1@gmail.com

ABSTRACT

The purpose of this study was to evaluate various types of fermentor for dry matter digestibility (DMD), organic matter digestibility (OMD), and the chemical composition of fermented seaweed. Five types of seaweed were used as substrates included green strain of *Kappaphycus alvarezii*, brown strain of *K. alvarezii*, *Gracilaria gigas*, *Sargassum* sp., and *Caulerpa* sp. The treatments were four fermentors, namely *Bacillus* sp. 2 mL/100 g of seaweed flour; 1.5% of tape yeast as a source of *Rhizopus* sp.; 1.5% of baker's yeast as a source of *Saccharomyces* sp.; a mix of *Bacillus* sp., tape yeast of *Rhizopus* sp. and baker's yeast of *Saccharomyces* sp. with compositions of 1 mL+1 g+1 g/100 g of seaweed flour; and control treatment. The results showed an increase in the percentage of DMD (21.94–27.76%) and OMD (8.35–11.66%) of all seaweed fermented using fermentor compared to control (DMD of 17.65–20.36% and OMD of 4.36–5.98%). Moreover, the highest result was obtained by the fermentor mix (DMD of 24.86–27.76% and OMD of 10.02–11.66%). Similar result was also found in the chemical composition of fermented seaweed, there was increase in protein content of 9.23–15.93% and nitrogen free extract (NFE) of 56.05–70.26% in each seaweed treated with fermentation using fermentors, compared to controls (protein of 8.82–11.54% and NFE of 52.26–65.72%). Furthermore, the highest value was shown by seaweed fermented with mixed fermentors (protein of 9.92–15.93% and NFE of 58.47–70.26%). Yet, the opposite result was present in the ash, crude fiber, and fat content of seaweed fermented using fermentors of which the lowest value was found in treatment of mixed fermentor.

Keywords: fermentation, fermentor, seaweed, quality, feed ingredients

ABSTRAK

Tujuan penelitian ini adalah mengevaluasi berbagai jenis fermentor terhadap pencernaan bahan kering (KBK), pencernaan bahan organik (KBO), dan komposisi kimia rumput laut terfermentasi. Lima jenis rumput laut digunakan sebagai substrat, yaitu *Kappaphycus alvarezii* strain hijau, *K. alvarezii* strain coklat, *Gracilaria gigas*, *Sargassum* sp., dan *Caulerpa* sp. Perlakuan yang diuji empat fermentor, yaitu *Bacillus* sp. 2 mL/100 g tepung rumput laut; 1,5% ragi tape sebagai sumber *Rhizopus* sp.; 1,5% ragi roti sebagai sumber *Saccharomyces* sp.; campuran *Bacillus* sp., ragi tape *Rhizopus* sp., dan ragi roti *Saccharomyces* sp. dengan komposisi 1 mL+1 g+1 g/100 g tepung rumput laut; serta kontrol. Hasil penelitian menunjukkan terjadi peningkatan persentase KBK (21,94–27,76%) dan KBO (8,35–11,66%) semua jenis rumput laut difermentasi dengan fermentor dibandingkan kontrol (KBK 17,65–20,36, dan KBO 4,36–5,98%) dan yang tertinggi dengan fermentor campuran (KBK 24,86–27,76 dan KBO 10,02–11,66%). Begitu juga yang terjadi pada komposisi kimia rumput laut difermentasi, terjadi peningkatan kadar protein (9,23–15,93%) dan bahan ekstrak tanpa nitrogen/BETN (56,05–70,26%) pada setiap rumput laut yang diberi perlakuan fermentasi menggunakan fermentor, dibandingkan kontrol (protein 8,82–11,54% dan BETN 52,26–65,72%), yang tertinggi ditunjukkan oleh rumput laut yang difermentasi dengan fermentor campuran (protein 9,92–15,93% dan BETN 58,47–70,26%). Namun, terjadi sebaliknya pada kadar abu, serat kasar, dan lemak rumput laut yang difermentasi dengan fermentor lebih rendah dibandingkan kontrol, dan terendah dengan perlakuan fermentor campuran.

Kata kunci: fermentasi, fermentor, rumput laut, kualitas, bahan pakan

INTRODUCTION

One of fishery products that can be used as a source of carbohydrates and also functions as a binder in fish feed is seaweed. Seaweed is listed as the commodity mostly produced in Indonesia. The production reached 54.12% or 1.7 million tons of total aquaculture production in 2013 of which 630.7 thousand tons was from South Sulawesi (Ministry of Marine Affairs and Fisheries Republic of Indonesia, 2014). In term of chemical composition, seaweed generally consists of water (27.8%), protein (5.4%), carbohydrates (33.3%), fat (8.6%), crude fiber (3%), and ash (22.25%) (Alamsyah, 2013). Moreover, seaweed also contains enzyme, nucleic acid, amino acid, vitamin (A, B, C, D, E, and K), and macro minerals such as nitrogen, calcium and selenium and trace minerals such as iron, magnesium, and sodium. The content of amino acid, vitamin, and mineral in seaweed is about 10 to 20 times higher if compared with land plants (Suparmi & Sahri, 2009). Seaweed which has been widely known and used by human consists of three classes, namely type of red (*Rhodophyceae*), brown (*Phaeophyceae*) and green (*Chlorophyceae*). Seaweed is also distinguished by the content of colloids. Seaweed group producing agar or commonly known as agarofit is namely *Gracilaria* and *Gelidium*, while the group producing carrageenan or termed as karaginofit consists of *Euchema* and *Kappaphycus*. Other group called alginofit which is able to produce alginate includes *Sargassum* and *Turbinaria*.

However, seaweed is a plant-based material that has cell walls and crude fiber. As stated by Utomo *et al.* (2011), the main obstacle in the utilization of plant-based materials as raw material for fish feed is the high content of crude fiber and the presence of anti-nutritional substances and amino acid composition those are different from the raw materials of animal protein. Excessive concentration of seaweed in feed results in high level of fiber whereas fish have limitations in digesting fiber. Therefore, concerning the feed formulation, seaweed should be at an optimum concentration.

One effort to improve the digestibility of feed can be applied by fermenting seaweed, which will be used as raw material for feed, using biodegradator. Fermentation is a chemical process of organic substrate change in the presence of biochemical catalysts, those are enzymes produced by certain microbes, with

the aim to simplify nutrient bond that is difficult to digest such as crude fiber which is found in fish feed ingredients. Seaweed fermentation can be conducted using microorganisms which function as biodegradator to improve the quality of feed ingredients. Utilization of *Rhizopus* sp. as fermentor to improve the quality of feed raw materials was reported by Amri (2007) in palm kernel meal and by Handajani (2011) in *Azolla* (aquatic plant) flour. Moreover, Amar *et al.* (2006) used *Bacillus* sp. to improve the quality of shrimp shells flour and Sukada *et al.* (2007) utilized *Saccharomyces cerevisiae* as fermentor in pollard, soybean bran, and cocoa shell. The purpose of this study was to evaluate the use of various fermentor types on dry matter digestibility (DMD), organic matter digestibility (OMD), and chemical composition of fermented seaweed.

MATERIALS AND METHODS

Materials used as substrates in fermentation process were five types of seaweed collected from seaweed farmers in Punaga Village, Takalar Regency. The five types of seaweed were *K. alvarezii* green strain, *K. alvarezii* brown strain, *Gracilaria gigas*, *Sargassum* sp., and *Caulerpa* sp. which had been cleaned and laid in the sun to dry. Furthermore, various types of seaweed were crushed and sieved up into flour.

Four treatments of fermentor were *Bacillus* sp. 2 mL/100 g of seaweed flour; 1.5% of tape yeast as a source of *Rhizopus* sp.; 1.5% of baker's yeast as a source of *Saccharomyces* sp.; a mix of *Bacillus* sp., yeast of *Rhizopus* sp. and baker's yeast of *Saccharomyces* sp. with compositions of 1 mL+1 g+1 g/100 g of seaweed flour; and control treatment. Three replications were performed for each treatment. *Bacillus* sp. used was the collection of the Laboratory of Nutrition and Food Technology, Faculty of Marine Sciences and Fisheries, Hasanuddin University with population density of 1.5×10^8 cfu/mL, tape yeast contained 1.6×10^7 cfu/g of *Rhizopus* sp., and baker's yeast contained 1.2×10^8 cfu/mL of *Saccharomyces* sp.

Fermentation was started by weighing each 100 g of seaweed flour which was later put in sealed plastic. Each 100 g of seaweed was treated by fermentor that has been dissolved in 20 mL of molasses and sprayed evenly using sprayer. Plastic wrapped the seaweed powder was sealed and the flour was incubated for 72 hours to give a chance to the fermentor to break the substrate apart. After 72 hours, the seaweed flour was

steamed in boiling water for 1–2 min to inactivate the fermentor. Furthermore, analysis of several variables was performed.

The parameters measured were dry matter digestibility (DMD) and organic matter digestibility (OMD), as well as the chemical composition of the seaweed included protein, fat, ash, crude fiber, and NFE through the proximate analysis. Formula for the determination of dry matter and organic matter digestibility at *in vitro* are as follows:

$$\text{Dry matter digestibility (DMD)} = \frac{\text{DM}_{\text{is}} - (\text{DM}_{\text{r}} - \text{DM}_{\text{b}}) \times 100}{\text{DM}_{\text{is}}}$$

$$\text{Organic matter digestibility (OMD)} = \frac{\text{OM}_{\text{is}} - (\text{OM}_{\text{r}} - \text{OM}_{\text{b}}) \times 100}{\text{OM}_{\text{is}}}$$

Note:

DM = dry matter (%)

OM = organic matter (%)

is = initial sample

r = residu

b = blanco

All observation data obtained in this study were analyzed descriptively by comparing the results obtained among treatments and also by comparing it with the supporting literature.

RESULTS AND DISCUSSION

The percentage of DMD produced in this study ranged from 17.65 to 27.76% while the percentage of OMD ranged from 4.36 to 11.66% (Table 1). The lowest percentage value of DMD and OMD was produced by various seaweeds of control treatment (without fermentors) and followed by treatment of single fermentor. Moreover, mix treatment of *Bacillus* sp., *Rhizopus* sp., and *Saccharomyces* sp. resulted in the highest value.

The high digestibility in the treatment of seaweed fermentation using fermentor of mixed *Bacillus* sp., *Rhizopus* sp., and *Saccharomyces* sp. was presumably due to the contribution of different extracellular enzymes released by each microorganism. According to Chu (2007), *Bacillus* sp. is a microorganism producing alkaline extracellular protease enzyme with a maximum proteolytic activity of 2,560 U/mL at temperature of 50 °C for 48 hours under optimal conditions. Alkaline protease is an enzyme that

hydrolyzes protein substrate in alkaline condition. *Rhizopus* sp. is a good producer of amylolytic enzyme, particularly glucoamylase (GA). This enzyme can lower both amylose and amylopectin by hydrolyzing α -1,4 and α -1,6 link of both glucoside starches and produce glucose (Nahar *et al.*, 2008). Jaelani *et al.* (2014) reported that the contribution of microorganisms in a fermentation can reduce the concentration of neutral detergent soluble, acid detergent soluble, hemicellulose, and lignin, as well as increase the protein content. According to Okone *et al.* (2007), degradation activity of fiber components by microorganisms involved in the fermentation process occurs during ensilage. Moreover, bacteria will convert non-fiber carbohydrates into organic acids (acetate, lactate, propionate, and butyrate) during ensilage. Thus, the final product is more digestible than the material without fermentation.

Digestibility is the first indication of the availability of various nutrients contained in certain feed ingredients for fish to consume. High digestibility reflects the contribution of a particular nutrient, while feed having lower digestibility indicates that the feed is less capable in supplying nutrients for a basic living or for the purpose of fish production. Muhtarudin and Liman (2006) found that higher DMD means an increase in OMD and higher utilization possibility of nutrients for the production, vice versa.

This study resulted in percentage value of DMD and OMD which was relatively similar to the research result of Hardana *et al.* (2013) by fermenting cocoa shells using *Aspergillus niger* of which the value of DMD ranged from 18.6% to 25.1% and OMD ranged from 3.4–13%. However, the result of this study was lower than that of research conducted by Afrijon (2011) on cocoa shell without urea treatment (DMD of 46.37%) and with 6% urea-ammoniation treatment (DMD of 52.80%) and was still much lower when compared to the research result of Yulistiani *et al.* (2012) which used different substrates, namely corn cobs treated with urea, *Aspergillus niger* and *Aspergillus niger* + 0.5 urea which produced higher DMD of 59.7%; 47.2%; and 50.9%, respectively. Tang *et al.* (2008) reported that OMD in feed supplemented with *Saccharomyces cerevisiae* reached a level of 56.8%.

The level of organic matter digestibility is relatively higher than the digestibility of dry matter. This is because the dry ingredients still contain ash, while the organic matter does not contain ash; thus, material without ash content is

Table 1. Percentage of dry matter digestibility (DMD) and organic matter digestibility (OMD) of seaweed with various fermentor treatments

Treatment	Parameter (%)	
	Dry matter digestibility (DMD)	Organic matter digestibility (OMD)
KAH.K	18.94 ± 1.02	5.98 ± 0.88
KAH.R	25.69 ± 0.82	9.65 ± 1.16
KAH.S	25.21 ± 1.24	9.25 ± 1.35
KAH.B	22.69 ± 1.35	11.12 ± 1.23
KAH.C	27.06 ± 1.40	11.54 ± 1.30
KAC.K	19.85 ± 1.16	5.49 ± 1.06
KAC.R	24.67 ± 1.22	9.71 ± 1.32
KAC.S	25.36 ± 1.37	9.46 ± 1.24
KAC.B	23.22 ± 0.86	11.57 ± 1.24
KAC.C	26.87 ± 0.62	11.66 ± 1.42
GG.K	18.93 ± 1.22	4.36 ± 1.28
GG.R	25.43 ± 1.15	9.04 ± 1.33
GG.S	25.78 ± 1.32	8.35 ± 1.08
GG.B	21.94 ± 1.27	9.43 ± 1.15
GG.C	27.76 ± 1.04	10.02 ± 0.7
SS.K	20.36 ± 1.08	4.87 ± 1.32
SS.R	26.35 ± 1.30	9.24 ± 1.36
SS.S	25.34 ± 1.26	9.15 ± 0.78
SS.B	24.36 ± 1.14	9.97 ± 0.89
SS.C	27.00 ± 1.18	10.59 ± 1.31
CS.K	17.65 ± 1.35	5.65 ± 1.02
CS.R	23.58 ± 1.15	8.97 ± 1.40
CS.S	23.14 ± 1.40	8.58 ± 1.32
CS.B	22.25 ± 1.27	9.36 ± 1.27
CS.C	24.86 ± 1.23	11.25 ± 1.25

Note: KAH = *Kappaphycus alvarezii* green strain; KAC = *K. alvarezii* brown strain; GG = *Gracilaria gigas*; SS = *Sargassum* sp.; CS = *Caulerpa* sp.; K = control; R = *Rhizopus* sp.; S = *Saccharomyces* sp.; B = *Bacillus* sp.; C = mix.

relatively easier to digest. As proven by Fathul and Wajizah (2010), the addition of Mn 40 mg/L and Cu 10 mg/L can increase the ash content which later may slow or inhibit the digestibility of dry matter of feed ingredients. In addition, organic matter is composed of carbohydrates, proteins, fats, and vitamins. Since organic matter is a part of dry matter, increase in the dry matter will also increase organic matter, vice versa. A similar result will also apply to the digestibility value, that is an increase in dry matter digestibility leads to increase in organic matter digestibility.

Results of proximate analysis of seaweed fermented with various fermentors (Table 2) showed that there was increased level of protein and NFE in each seaweed treated with fermentation using fermentors, compared with control. Furthermore, the highest value was found in seaweed fermented with a mix of *Bacillus* sp., *Rhizopus* sp., and *Saccharomyces* sp. However, the content of ash, crude fiber, and fat in seaweed fermented with fermentors was lower than that of control, and the lowest result was obtained in the treatment of mixed fermentors.

Table 2. Result of proximate analysis of seaweed by various fermentors

Treatment	Parameter (%)				
	Ash	Protein	Fat	Crude fiber	NFE
KAH.K	8.95±0.78	11.54±1.27	1.64±0.03	15.73±1.18	62.14±1.35
KAH.R	8.49±0.54	13.13±1.05	0.81±0.01	11.37±1.04	66.19±1.16
KAH.S	7.48±0.89	13.37±1.12	1.30±0.04	11.70±1.13	66.15±1.22
KAH.B	8.08±0.50	14.06±1.34	0.81±0.01	11.06±1.26	65.99±1.08
KAH.C	6.39±0.78	15.93±1.21	0.20±0.05	11.23±1.24	66.25±1.14
KAC.K	9.00±1.02	11.27±1.4	1.74±0.02	14.90±1.35	63.09±0.89
KAC.R	6.79±0.76	14.32±1.22	0.82±0.05	11.56±0.89	66.50±1.65
KAC.S	6.95±0.45	14.31±1.35	0.84±0.03	11.39±1.33	66.51±1.25
KAC.B	7.02±0.82	14.21±1.3	0.83±0.02	11.26±1.14	66.69±1.27
KAC.C	5.88±0.35	15.27±1.27	0.83±0.05	10.73±1.12	67.30±1.22
GG.K	10.14±0.98	8.82±0.8	0.91±0.01	15.56±1.27	64.58±1.14
GG.R	9.92±0.63	9.61±1.05	0.71±0.03	11.04±1.06	68.73±0.67
GG.S	10.97±0.75	9.23±0.96	0.29±0.01	11.83±1.01	67.67±1.27
GG.B	10.28±0.86	9.42±1.27	0.73±0.04	12.47±0.85	67.10±1.35
GG.C	9.48±0.78	9.92±1.03	0.82±0.04	10.83±1.22	68.95±1.16
SS.K	11.18±0.65	10.88±1.09	1.25±0.03	10.97±1.08	65.72±1.42
SS.R	9.07±0.45	11.46±0.69	1.00±0.02	7.91±1.25	70.55±0.76
SS.S	9.06±0.48	12.95±1.07	0.33±0.02	9.37±1.07	68.29±1.05
SS.B	9.66±0.94	13.07±1.30	0.85±0.03	8.35±0.59	68.07±1.20
SS.C	6.68±0.65	13.55±1.23	0.75±0.05	8.76±1.27	70.26±1.15
CS.K	13.19±1.4	10.56±1.28	1.91±0.03	22.08±1.60	52.26±1.27
CS.R	9.00±0.87	11.04±1.04	1.38±0.05	20.53±1.02	58.05±1.08
CS.S	7.22±0.99	12.69±1.08	1.22±0.05	20.27±1.24	58.60±1.13
CS.B	7.21±0.75	12.82 ± 0.88	1.53±0.01	20.20±1.07	58.24±1.28
CS.C	7.29±0.68	12.96 ± 1.24	1.31±0.03	19.97±1.21	58.47±0.92

Note: KAH = *Kappaphycus alvarezii* green strain; KAC = *K. alvarezii* brown strain; GG = *Gracilaria gigas*; SS = *Sargassum* sp.; CS = *Caulerpa* sp.; K = control; R = *Rhizopus* sp.; S = *Saccharomyces* sp.; B = *Bacillus* sp.; C = mix.

Increased level of protein and NFE of seaweed fermented with various fermentors was caused by enzymatic by fermentor and the addition of protein in fermentor cell. According to Sukaryana *et al.* (2011), work result of enzyme produced by microorganisms leads to changes in the fermentation process (either in aerobic or anaerobic condition). Akinfemi *et al.* (2009) showed that increase in crude protein content on fermentation of solid substrate occurs as a result of hydrolysis of starch into sugar when fungus degrades and dissolves the substrate which later will be used by the fungus as a source of

carbohydrates to synthesize fungus biomass that is rich in protein. Further, fermentation of solid substrate on lignocellulosic feed ingredients using microorganisms can increase the protein content and lower the fiber content (cellulose and hemicellulose, neutral detergent soluble, acid detergent soluble). Jalil *et al.* (2015) reported that sago fermented with *R. oligosporus* could increase the reducing sugar by 61%, glucose and fructose by more than 100%, soluble protein by 1% and amino acids by 1.5% to 38.2%.

It is known that crude fiber content in feed ingredients affect the digestibility or degradation

of dry matter and organic matter. According to Suprpto *et al.* (2013), crude fiber content is not only able to affect the digestibility of dry and organic matter but also may affect the digestibility of other compounds. Rostika and Safitri (2012) stated that fermentation can improve the nutritional quality of feed ingredients through fiber degradation, fat degradation, and an increase in crude protein, which further will increase the digestibility of feed. Oduguwa *et al.* (2008) reported that fermentation with fungus *R. oligosporus* on rice bran and coffee hull increased NFE and lower fiber content. Mulia *et al.* (2015) also reported that tofu waste fermented with *R. oligosporus* successfully decreased crude fiber content by 38–40%.

A decrease in crude fiber was possibly caused by the breaking down of complex substances in the substrate into simpler compounds by fungus *T. harzianum*. An increase in protein was sourced from biomass of growing fungus which is also a single cell protein (SCP). Biomass of fungus contained in the fermented material can contribute as a source of protein and is also expected to enhance the immune response of fish and shrimp (Kang *et al.*, 2010). Compared with the crude protein content of corn cobs without treatment (4.33%), crude protein content in corn cobs biodegraded using fungus *Pleurotus ostreatus* and treated with the addition of urea increased twice (Hatta & Sugiarto, 2015). It is also explained that crude protein increase in fermented young corn cob can also occur due to the growth and proliferation of fungi during fermentation. In general, all end products of fermentation usually contain simpler and easier compounds to digest than the original materials; thus, the nutritional value of the ingredients will increase. Various types of microorganisms have the ability to convert starch into proteins by the addition of inorganic nitrogen through fermentation.

CONCLUSION

Fermentation of seaweed *Kappaphycus alvarezii* green strain, *K. alvarezii* brown strain, *Gracilaria gigas*, *Sargassum* sp., and *Caulerpa* sp. using fermentors of *Bacillus* sp. 2 mL/100 g of seaweed flour; 1.5% of ragi tape yeast as a source of *Rhizopus* sp.; 1.5% of baker's yeast as a source of *Saccharomyces* sp.; mixed fermentors (*Bacillus* sp., tape yeast of *Rhizopus* sp. and baker's yeast of *Saccharomyces* sp. with compositions of 1 mL+1

g+1 g/100 g of seaweed flour) could increase the percentage of dry matter digestibility (DMD) and organic matter digestibility (OMD), protein content, and nitrogen-free extract (NFE), as well as decrease the content of crude fiber, fat, and ash of seaweed. The best fermentor was the mix of *Bacillus* sp., tape yeast *Rhizopus* sp., and Baker's yeast *Saccharomyces* sp.

ACKNOWLEDGEMENTS

The author would like to thank the Directorate General of Higher Education for funding this research through the National Priorities Research Master Plan for the Acceleration and Expansion of Indonesian Economic Development 2011–2025 (Penelitian Prioritas Nasional Masterplan Percepatan dan Perluasan Pembangunan Ekonomi Indonesia, PENPRINAS MP3EI 2011–2025) of The Fiscal Year 2015.

REFERENCES

- Afrijon. 2011. Influence of using urea in pod cacao amination for dry matter and organic digestibility by *in vitro*. *Jurnal Embrio* 4: 1–5.
- Akinfemi A, Adu OA, Doherty F. 2009. Assessment of the nutritive value of fungi treated maize cob using *in vitro* gas production technique. *Livestock Research for Rural Development* 21: 188.
- Alamsjah MA. 2013. *Gracilaria* sp. waste, *Lactobacillus* sp. and *Chlorella* sp. integration on intensive aquaculture with aquaponic system. *Journal of Natural Sciences Research* 3 (11): 66–77.
- Amar B, Philip R, Bright Singh IS. 2006. Efficacy of fermented prawn shell waste as a feed ingredient for indian white prawn, *Fenneropenaeus indicus*. *Aquaculture Nutrition* 12: 433–442.
- Amri M. 2007. Effect fermented palm kernel cage portion in feed of common carp *Cyprinus carpio* L. *Jurnal Ilmu-Ilmu Pertanian Indonesia* 9: 71–76.
- Chu WH. 2007. Optimization of extracellular alkaline protease production from species of *Bacillus*. *Journal of Industrial Microbiology and Biotechnology* 34: 241–245.
- Fathul F, Wajizah S. 2010. Additional micromineral Mn and Cu in ration to rumen biofermentation activities of sheep *in vitro* method. *Jurnal Ilmu Ternak dan Veteriner* 15: 9–15.

- Handajani H. 2011. Optimalisasi substitusi tepung *Azolla* terfermentasi pada pakan ikan untuk meningkatkan produktivitas ikan nila gift. *Jurnal Teknik Industri* 12: 177–181.
- Hardana NE, Suparwi, Suhartati FM. 2013. Cocoa pods (*Theobroma cacao* L.) fermentation using “probiotik x” its effect on dry matter digestibilities (DMD) and organic matter digestibilities (OMD) *in vitro*. *Jurnal Ilmiah Peternakan* 1: 781–788.
- Hatta U, Sugiarto. 2015. Produksi tepung tongkol jagung muda hasil biodegradasi kapang *Pleurotus ostreatus* dengan enzim pemecah serat dan implikasinya pada pakan ayam pedaging. *Jurnal Ilmu-Ilmu Peternakan* 25: 1–7.
- Jaelani A, Djaya S, Rostini T. 2014. Characteristics and nutrition silage duckweed (family *Lemnaceae*) addition with different additives. *International Journal of Biosciences* 5: 144–150.
- Jalil AA, Abdullah N, Alimon AR, Abd-Aziz S. 2015. Nutrient enhancement of ground sago *Metroxylon sagu* Rottboll pith by solid state fermentation with *Rhizopus oligosporus* for poultry feed. *Journal of Food Research* 4: 1–15.
- Kang HY, Yang PY, Dominy WG, Lee CS. 2010. Bioprocessing papaya processing waste for potential aquaculture feed supplement—economic and nutrient analysis with shrimp feeding trial. *Bioresource Technology* 101: 7973–7979.
- Ministry of Marine Affairs and Fisheries Republic of Indonesia. 2014. Analisis data kelautan dan perikanan. Jakarta: Kementerian Kelautan dan Perikanan.
- Muhtarudin, Liman. 2006. Determination of utilization level of organic mineral to improve rumen bioprocess of goat by *in vitro* method. *Jurnal Ilmu-Ilmu Pertanian Indonesia* 8: 132–140.
- Mulia DS, Yulyanti E, Maryanto H, Purbomartono C. 2015. Quality Improvement of tofu waste as the raw material of fish feed with fermentation of *Rhizopus oligosporus*. *Sainteks* 12: 10–20.
- Nahar S, Hossain F, Feroza B, Halim MA. 2008. Production of glucoamylase by *Rhizopus* sp. in liquid culture. *Pakistan Journal of Botany* 40: 1693–1698.
- Oduguwa OO, Edema MO, Ayeni AO. 2008. Physico-chemical and microbiological analyses of fermented corncob, rice bran and cowpea husk for use in composite rabbit feed. *Bioresource Technology* 99: 1816–1820.
- Okone A, Yimamu A, Hanada M, Izumitta M, Zinong M, Okamoto M. 2007. Ensiling characteristic of daikon *Raphana sativum* by product and its potentials as an animal feed resource. *Journal Animal. Feed Science and Technology* 136: 248–264.
- Rostika R, Safitri R. 2012. Influence of fish feed containing corn-cob was fermented by *Trichoderma* sp., *Aspergillus* sp., *Rhizopus Oligosporus* to the rate of growth of Java barb *Puntius Gonionitus*. *APCBEE Procedia* 2: 148–152.
- Sukada IK, Biduri IGNG, Warmadewi DA. 2007. The effect of fermented pollard, soybean hull, and cocoapod with “yeast culture” on carcass weight and meat cholesterol of male Bali duck. *Majalah Ilmiah Peternakan* 10: 1–10.
- Sukaryana Y, Atmomarsono U, Yunianto VD, Supriyatna E. 2011. Improvement of crude protein and crude fiber digestibility of fermented product of palm kernel cake and rice bran mixture for broiler. *Jurnal Ilmu Ternak dan Veteriner* 1: 167–172.
- Suparmi, Sahri A. 2009. Mengenal potensi rumput laut: Kajian pemanfaatan sumber daya rumput laut dari aspek industri dan kesehatan. *Majalah Ilmiah Sultan Agung* 44: 95–116.
- Suprpto H, Suhartati FM, Widiyastuti T. 2013. Digestibility of crude fiber and crude fat complete feed jute waste with different protein sources on post weaning etawa cross breed goat. *Jurnal Ilmiah Peternakan* 1: 938–946.
- Tang SX, Tayo GO, Tan ZL, Sun ZH, Shen LX, Zhou CS, Xiao WJ, Ren GP, Han XF, Shen SB. 2008. Effects of yeast culture and fibrolytic enzyme supplementation on *in vitro* fermentation characteristics of low-quality cereal straws. *Journal of Animal Science* 86: 1164–1172.
- Utomo NBP, Nurfadhilah, Ekasari J. 2011. Fermentation of *Azolla* sp. leaves and the utilization as a feed ingredient of tilapia *Oreochromis* sp. *Jurnal Akuakultur Indonesia* 10: 137–143.
- Yulistiani D, Puastuti W, Wina E, Supriati. 2012. Effect of processing on nutritive value of corn cobs: chemical composition and *in vitro* digestibility. *Jurnal Ilmu Ternak dan Veteriner* 17: 59–66.