

METHODOLOGICAL APPROACHES IN RUMINANT METABOLIC RESEARCH¹

PENDEKATAN METODOLOGI DALAM PENELITIAN METABOLISME PADA RUMINANSIA

Djokowoerjo Sastradipradja²

ABSTRACT

Climatic **factors** of high temperatures and humidities of humid tropical countries, alongside with the **occurrence** of diseases and parasites have created a complex situation in terms of the effect on animal production. Studies on quantitative relationships for better understanding of digestion, metabolism and related areas are lacking. There are however a number of appropriate methodologies available for conducting **onfarm** metabolic research aiming at giving better understanding of animal production and health in the humid tropics. The paper discusses methodologies and experimental approaches developed in the authors's laboratory on growing, **draft**, pregnant and lactating **ruminants**. Several examples of the results are presented.

ABSTRAK

Faktor-faktor suhu tinggi dan kelembaban lingkungan tropika lembab, digabungkan dengan tingginya kejadian penyakit dan parasit menciptakan situasi yang kompleks yang berpengaruh negatif terhadap upaya produksi ternak. Penelitian-penelitian mengenai aspek kuantitatif fungsi pencernaan, metabolisme hewan dan bidang-bidang terkait boleh dikatakan sangat minim. Meskipun demikian, sebenarnya cukup tersedia metodologi yang sesuai untuk melaksanakan "onfarm metabolic research" yang bertujuan memperoleh pemahaman yang mendalam mengenai produksi dan kesehatan ternak di daerah tropika lembab. Makalah ini membahas beberapa metodologi dan pendekatan percobaan yang dikembangkan di laboratorium penulis pada hewan ruminansia yang sedang tumbuh, bunting, laktasi dan ternak kerja. Beberapa contoh hasil penelitian disajikan untuk ilustrasi.

¹ Center for Metabolic Research, presented at a one-day seminar "Improvement of Animal Performance through Onfarm Metabolic Research" organized by the Faculty of Veterinary Medicine Bogor Agricultural University (IPB) at JL. Taman Kencana 3 Bogor 16151, 30 April 1997, INDONESIA

² Department of Physiology and Pharmacology, Faculty of Veterinary Medicine, Institut Pertanian Bogor (Bogor Agricultural University), Jalan Taman Kencana 3, Bogor 16151, INDONESIA

INTRODUCTION

Food is necessary to built up tissue and to act as a source of energy. Food has to be digested into absorbable units in the gastrointestinal tract. Metabolism is the process refered to all the chemical and energy transformations that occur in the body.

The humid tropical countries have special and often unique problems associated with animal production. Both imported and indigenous animals used to build the livestock industry shows variation in adaptability to the existing **agro-ecological** setting of the region. The countries have dietary materials available, particularly for ruminants, which in **many** instances have not been evaluated for their ability to support the desired production. Humid tropical countries have adverse climatic factors of high temperatures and humidities and, in addition, there are a wide variety of diseases and parasites which reduce production. Direct effects of these factors **and** interactions between the factors have created a complex situation in terms of the effect on production.

As qualitative knowledge increased, detailed consideration is given to develop quantitative relationships to increase further understanding and integrate various aspects, **e.g.** to bring together quantitative approaches concerned with elucidating mechanisms, used in the **study** of digestion (monogastric and ruminant), metabolism and related areas. In his address to the 11th Symposium on Energy Metabolism of **Farm** Animals on "Past Achievements and Future Perspective in Energy Metabolism", the late Sir Kenneth L. **Blaxter** (1989) made the following **remarks**: *"Current world literature did not contain much information which relates to climatic and seasonal or exogenous hormonal effects on metabolism, pregnancy in mammals, egg secretion in birds and effects of muscular work have not been considered. Nor has the comparative aspect of metabolism been considered On the practical side we need these are items of commerce. We also need, more accurate and rapid methods for estimating the energy values of fee & sice; to take into account breed and strain differences in' animal requirements. Better estimates of the composition of body gains (retentions) are required. New demands will arise as a result of modern techniques of genetic manipulation of livestock and the use of exogenous hormones derived through, biotechnology, to increase reproductive performance and the rate and composition of growth. On the fundamental side, we must explore aspects of adaptation of metabolism to undernutrition, season and climate at the cell, organ and whole animal levels and unravel the complexity of the endocrinee and neural control mechanisms involved focussing on*

the roles of the sympathetic nervous system and growth factors. New techniques must be explored since they may well enable studies to be made in natural environments rather than the laboratory. Biochemical methods of assessing the energy status of animals demands attention". These hold true for the humid tropical situation as well.

Brody (1945) stated that growth is the basis of and closely related to many animal productive processes, which includes egg, **milk**, fat and other production. He defined growth as the constructive and assimilatory synthesis of one substance at the expense of another, **i.e.** nutrient which undergoes dissimilation. Growth, and hence animal production, is biological synthesis, production of new biochemical units and it is the aspect of development concerned with the increase in living substance or protoplasm and includes one or all of the following processes, **e.g.** cell multiplication, cell enlargement, and incorporation of material taken from the environment. From the practical point of view of quantitative measurement of growth of the organism as a whole, nonprotoplasmic inclusions in the body must be considered as parts of the growth process.

The aforesaid processes are more dramatic during periods following starvation and injury. Seasonal variations in animal feed supplies **often** cause interrupted patterns of growth in young animals due to feed restriction resulting from seasonal variations or other causes. A subsequent restoration of feed supplies causes the animals to exhibit enhanced gain in weight called compensatory growth. The underlying physiological processes of growth are complex involving feed utilization, visceral tissue mass and **carcass** protein and fat deposition, variation in energy requirements according to a particular condition and also in other nutrients, to meet maintenance and body performance, partition of nutrients, endocrine control, etc. So **defined**, growth (which includes production) is inseparable from metabolism.

Our studies on the responses of ruminant (and other livestock) animals to different rations or related treatments rely basically on traditional balance methods alongwith proximate analysis of foodstuffs and waste. To **measure** the transfer of substances within the intact animal and their partition to organs and tissues, tracer isotope methodologies are used in combination with experimental approaches such as the application of the Fick principle and dilution techniques. The data range over a wide field area and include important considerations as the energy cost of maintaining animals, of transforming feed into body tissue and the desired animal products. Data on body composition enable assessment of growth responses to changes in nutrition or in the

environment. The studies are reviewed in this presentation highlighting on experience in the use of methodologies appropriate to the existing conditions.

The methodologies and experimental approaches presented are based on the work on growing animals. As the discussions proceed to touch on other **animal productive** states, the appropriate methodologies will be inserted.

GROWING RUMINANTS

Animal Balance Trial

The balance trial provides information related to the requirements of nutrients for growth, maintenance and production and the **availability** of nutrients from **feedstuffs**. It is used to show changes in percentage absorption of nutrients as a **function** of the level of nutrients in the diet. For energy balance, the energy categories of feed, **i.e.** gross energy (GE), faecal energy (**FE**), digestible energy (DE), urinary energy (UE), Energy loss with digestive gas production (**En-CH₄**), metabolizable energy (ME), energy expenditure (EE) = health production (HP) and retained energy (RE), should be calculated. Energies in feed, **faeces** and urine are determined by bomb calorimetry or calculated from organic **ingredients** multiplied by their respective caloric equivalents, **e.g.** UE may be found from $\text{g urinary-N} \cdot d^{-1} \times 34.0 \text{ kJ}$, but energy in gases and **CH₄** is rather difficult to measure. It could be reasonably well approximated from gas test results or simply taken as 8-10 % of GE, or using **Blaxter's equation** $\text{En-CH}_4 = 4.28 + 0.059 D \text{ Kcal/100Kcal GE}$ (D = digestibility of energy, %). For the energy balance, data on HP and RE are needed. When physical exercise is absent, it is safe to use the relationship $\text{ME} = \text{HP} + \text{RE}$. Since ME is found from GE **subtracted** by all energy **found** in waste, either HP or RE should be determined while the other could then be calculated by difference **from** the ME. If HP should be **measured**, a suitable technique of animal calorimetry is required. On the other hand if RE is chosen to be determined, the slaughter technique or a suitable **in vivo** body composition technique should be used. Alongside energy balance, protein (or N) balance could be simultaneously measured.

Whole Animal Calorimetry and Body Composition

All major components of the body are in a continuous **state of flux**: intakes, excretions, degradations and resynthesis. All **endproducts** of digestion not only act as substrates for the accretion of body tissue (anabolism), but also **serve** as **precursors** for the production of the high-energy compounds needed (catabolism) for maintenance and to provide the energy for the synthesis of the macromolecules of the body. In addition, there is an extensive **interconversion** of metabolites and the rate of productive processes in **the** body is governed by the complex interplay between the intermediary metabolism of the **various** nutrients **and** the **rate** of synthesis and degradation of tissue lipid, protein, nucleic acid and carbohydrate **stores**.

1, Measurement of Whole Body HP by Carbondioxide Entry Rate' Technique

Due to unavailability of respiration chamber **calorimetry** for large animals, but on the other hand we have in our **laboratory** at our disposal radioisotope detection equipment, Carbondioxide Entry Rate Technique (CERT) using tracer ¹⁴C-bicarbonate is the method of choice to measure whole body energy expenditure of small ruminants (Sastradipradja, 1992). The method involves primed continuous infusion of the label solution into a blood vein and after reaching steady state condition of isotope concentration in body fluid bicarbonate, serial blood samples are withdrawn. The CO₂ production rate (**rCO₂**) is calculated by dividing the rate of label infused by the plateau specific activity of blood bicarbonate. At an accepted RQ value, the **rCO₂** value is converted into its energy equivalent which is HP. Double polyethylene catheters are implanted in the jugular veins for minimal disturbance to the animal, easy delivery of isotope and convenient serial blood sampling. It is important to measure the specific activity of CO₂ over a sufficiently long period of time (**e.g.** 12h or longer) to ensure a more representative mean value of specific activity. For such long trials, instead of blood, samples of any body fluid **can** be taken (**e.g.** saliva instead of blood). To get faster attainment of stable specific activity we apply the primed-continuous infusion technique and our experience shows that the ratio between primer dose and infusion rate per minute is 80 to 1 giving satisfactory results. Primer dose is 1 ml of NaH¹⁴CO₃ = 40 **μCi** delivered within 1 min. followed by continuous infusion at 0.5 **μCi/min**. The CO₂ production **can** be estimated **from** the plateau specific activity according to the equation (Corbett *et al.*, 1971) :

$$\text{CO}_2 = \frac{\text{rate of tracer bicarbonate infused}}{\text{plateau specific activity of CO}_2}$$

The advantage of CERT is that the **animal** is **free** to move **around without restraint**. The use of CERT also enables **measurement** of **gluconeogenesis** involving $^{14}\text{CO}_2$ fixation. Another advantage of **CERT** is **the** ability of **measuring** glucose kinetics during the **same** trial by administering ^3H -glucose at **the same time** with the bicarbonate label.

2. *Measurement of HP as the Difference Between ME and RE*

In our work with **swamp** buffaloes during the last four years, we have calculated HP as the difference between ME and RE included physical work (**Mahardika et al.**, 1997) with satisfactory results (see exercise metabolism below).

3. *Heart Rate as A Predictor of HP*

We are still developing this technique on small **ruminants** and cattle following the success of **longterm heart** rate measurement in swamp buffaloes using the Polar Sport Tester (Finland).

4. *Estimation of Body Composition in vivo*

4.a *Urea Space Technique* (Rule et al., 1986)

Urea is used as a marker for body water. A measured amount of urea in saline is injected into the jugular vein within one minute. Exactly 12 minutes after the injection, a sample of blood is withdrawn from the jugular vein and the concentration of urea determined. Urea space is found by dividing the dose (mg) of urea **infused** by the increment of blood urea concentration following the **infusion** from the **preinfusion** value times body weight times 10. The empty body water (EBW, %) = $59.1 + 0.22 \text{ US (\%)} - 0.04 \text{ BW}$, while empty body fat (%) = $19.5 - 0.31 \text{ US (\%)} + 0.05 \text{ BW}$. The choice of this technique is necessary considering the hazards involved in the use of radioisotope labelled water on large ruminants **and** the costly slaughter technique with ruminant livestock animals. We have introduced this technique in our work with goats and sheep. Validation of the *in vivo* body composition method using US was separately done on four goats by doing slaughter technique (ST) analysis in addition to the *in vivo* calculations on the same animals in question (**Arta Putra et al.**, 1997). EBW estimated according to Rule et al. (1986) resulted in an underestimation of 4.2% from the EBW value by ST. Therefore, the use of US in goats required a correction factor of 1.044. For use with small ruminants, **Panaretto's** equation for body protein (**Panaretto**, 1963) and for body fat (Panaretto and Till, 1963) are

based on EBW. The use of our **corrected** values in aforesaid **equations** resulted in **good** agreement with our ST results. The **results** are **presented** in **Table 1**.

Table 1. **Body Composition (% BW) According to Slaughter Technique Analysis and in vivo Equations Estimated from Four Female Growing Goats**

	Body water		Body protein		Body fat	
	ST	Eq. 1	ST	Eq. 2	ST	Eq. 3
	----- %BW -----		----- %BW -----		----- %BW -----	
Mean ± SD	60.9±5.94	60.9±0.06	20.2±0.39	20.2±0.02	16.2±5.74	17.6±0.08

ST = slaughter technique analysis

Eq. 1 = EBW according to Rule *et al.* (1986) US equation multiplied by 1.044,

Eq. 2 = Panaretto (1963) body protein equation using EBW value from Eq. 1,

Eq. 3 = Panaretto and Till (1963) body fat equation using EBW value from Eq. 1.

Validation results with the Javanese thin tail (JIT) sheep (Saka and Sastradipradja, unpublished data) revealed that only the equation which relate EWB with US was significantly different from the regression equation according to Rule *et al.* (1986), as follows

$$EBW (\%) = 64.1 - 21.1US\% - 0.378BW$$

$$(P < 0.05; S_{y.x} = 2.185; R^2 = 0.425)$$

Consequently, only this equation can be used for sheep with correction factors respectively for the intercept, US and BW to be 1.0846, -95.9091 and 9.45. From our studies with female swamp buffaloes, the equation of Rule *et al.* (1986) if applied for working buffaloes should be modified to:

$$Fat = 13.69 - 0.21US + 0.03 BW; Protein = 19.7 + 0.08 US + 0.11 BW.$$

4.b Body Density Method (Kleiber, 1961)

Application of this technique in small ruminants and cattle is **being** conducted **after** it has been **successfully** practiced in the swamp buffalo. Contrary to the situation in swamp buffaloes who love wading in water, other livestock animals will resist to immersion in water. Therefore, a solid frame should be constructed where the animal can be immobilized and the whole device including the animal can be subsequently lowered in the water tank leaving the head above water level.

Supporting Measurements on Metabolism and Digestion

Glucose is an **important nutrient** for all tissues, especially it is indispensable as energy source for nerve cells and for foetus in pregnant animals, for the synthesis of lactose, fat and providing carbon skeleton for **others**. An important component of **growth** is fat synthesis. Glucose is required during growth, in particular for the supply of the NADPH via the pentosephosphate pathway for **fat** synthesis. **A** lower amount is required for **protein** synthesis. The major portion of glucose available to the **ruminant** is supplied by gluconeogenesis. Additional demands for **glucogenic precursors occur during** lactation, namely for the lactose secreted in **the** milk and for the reducing power needed **to** provide energy for milk synthesis. Impaired gluconeogenesis is believed to contribute to various metabolic disorders frequently seen in ruminants, such as acetonemia in dairy cows and pregnancy toxemia or twin lamb disease in sheep. In ruminant studies, glucose metabolism warrants attention.

Successful operation of whole animal metabolism requires low levels to function properly. Many metabolic peculiarities of ruminants stem from the proper functioning of the splanchnic region. It is understandable that special attention should be given to metabolism in this region.

A great deal of research has been carried out on the digestive system of the ruminant giving **understanding** of the characteristic metabolism which cope with the peculiar products of microbial digestion. We attempted to measure products of ruminal digestion and integrate it for our overall understanding of animal performance.

1. Gluconeogenesis Involving CO₂ Fixation

Tracer labeling of the bicarbonate pool (CERT) may enable one to estimate gluconeogenesis (GNG) **from** carbon transfer involving **CO₂** fixation into the glucose pool. The rationale involves using the transfer quotient (TQ) between **CO₂** and glucose to assess the extent of **CO₂** fixation in GNG from precursors like propionate and lactate. The TQ in question is calculated as the ratio between the plateau specific activities (per at. C) of **¹⁴C** in glucose (product) and in bicarbonate (precursor). The rate of GNG involving **CO₂** fixation is found by multiplying the value of the glucose **flux** by the TQ times 6. Measurement of the glucose **flux** and pool size is done simultaneously with the CERT infusion in the same animal by pulse injection with tritiated glucose (Sastradipradja, 1992).

2. Metabolism of the Portal Drained Viscera

Many metabolic peculiarities of ruminants stem **from** the viscera, that is a group of organs whose blood supply **drains** into the portal vein. Quantification of nutrients in the **splanchnic** region is useful to **understand the** metabolism in this region. We have experience to **study** this **aspect** in small ruminants by the **arteriovenous** difference technique (Katz and **Bergman**, 1969) involving surgical procedures to obtain **samples from** the **mesenteric, portal and** hepatic veins and of any artery (Astuti, 1995; Sastradipradja *et al.*, 1997). Portal blood flow were measured on unanaesthetized animals (**i.v.** injection of **xylazine** 0.05 **ml/kg** BW, followed by **i.v.** injection of ketamine 0.11 **ml/kg** BW). Lidocaine (1 **ml/** animal) was applied locally on the site of incision. The primer dose of **PAH-³H** was 5.75 **µCi** delivered in one minute via the jugular vein followed by continuous infusion at 0.6 **µCi/min** into the right mesenteric vein. After around two hours of **infusion**, blood samples were collected for analysis of blood gasses (**CO₂, O₂, VFA**) and other relevant constituents. Metabolism of the region and absorption rates of nutrients can be measured. The technique is invasive however, which need appropriate skills to perform. So far we haven't been successful in placing permanent catheters in aforesaid blood vessels. Feasible techniques to measure ruminal fermentation production rates would be an alternative approach to measure the nutrients' supply to the animal.

3. In vitro Techniques for Ruminal Fermentation

The techniques closely simulate **in vivo** conditions especially for **rumen** fermentation studies: rate of formation of endproducts (VFA), rate of release of **NH₃**, production rate of microbial protein and gas production.

3.a' *Microbial Protein Synthesis* (Suwandyastuti *et al.*, 1985)

A shortterm incubation of **rumen** contents at 39 °C in an artificial **rumen**: test **tubes**, glass syringes or **flasks**. The anaerobiosis is maintained by introducing **N₂** gas replacing the air in the artificial **rumen** above the incubation mixture. **³³P** is used as tracer for measuring microbial protein synthesis (MPS).

3.b *Incubation of Rumen Contents in Glass Syringes (gas test)* (Menke and Steingass, 1988)

The amount of gas which is released when feedstuffs are incubated **in vitro** with **rumen** fluid measures **CO₂** and **CH₄** production. It is closely related to digestibility and **consequently** to the energetic feed value of feedstuffs for ruminants. The incubation **mixture** consists of feed

sample, main and **trace** element solution, buffer solution, **rezazurin solution** (indicator), **reduction** solution and rumen fluid. Hundred ml *syringes* are used **preheated** at 39 °C. The syringes **plus** contents **free** of air bubbles are incubated in a (39 °C) preheated incubator. The **volume** of **gas** in the piston is read off eight hours after the **start** of the incubation (**V₈**). Incubation is **continued** and the final reading is done **after** 24 hours (**V₂₄**). Regression equations will be derived relating ME with gas production, feed nutrient components and **digestibilities**.

3.c VFA Production Rates: *Zerotime* in vitro *Method* (Whitelaw *et al.*, 1970)

A sample of **rumen** contents is taken and subsamples incubated *in vitro* under anaerobic **conditions**. The rate of production of individual and total VFA is calculated **from** the increments in acid concentration obtained by incubating the subsamples for different periods and extrapolating back to zero time incubation to give the rate of VFA production per unit volume at the time the sample was removed. Equations for performing the calculations are given by **Whitelaw et al.** (1970). The **rumen** volume should be known in order to calculate total ruminal production.

3.d *Use of Nylon Bags Incubated in The Rumen* (IAEA, 1985)

The digestibility test requires animals fitted with permanent **rumen** fistulae. Nylon bags should have a pore size 20 - 40 mm, dimensions 15 x 8 cm, sample size 3-5 g of air-dry feed, ground through screen 2-5 mm, incubation times up to 24h for protein concentrates, up to 72h for roughage feeds. The technique can describe both the rate and the extent of degradation, affected by the **rumen** environment such as ammonia level, pH, **type** of feeds, trace minerals, etc.

4. In vivo *Estimation of Digestion*

Evaluation on the intake **and** digestion characteristics of feedstuffs, needs quantitative data to describe the movement of **digesta** along the gastrointestinal tract.

4.a *Estimation of Rumen Volume* (IAEA, 1985)

The volume of **rumen** liquid is **estimated from** the dilution of ⁵¹Cr-EDTA tracer introduced **intraruminally** by way of a syringe and needle in a period between **rumen** contractions. After 1 hour and then every hour up to eight hours, **sample/withdraw** representative **rumen** contents by **way** of a stomach tube. The animal should receive a (nearly) continuous feeding regime. Plot

log concentration against time and find the intercept to give the ^{51}Cr concentration at zero time. Rumen volume is calculated as dose divided by zero time concentration. Flow of rumen content is obtained from the kinetics' data. $^{51}\text{CrCl}_3$, obtained from Amersham UK, mixed with Na-EDTA will easily form $^{51}\text{Cr-EDTA}$. Alternatively, the volume of rumen can also be estimated from *per oral* administration of an aliquot of labeled water ($^3\text{H}_2\text{O}$ or D_2O) and follow the tracer disappearance with time in rumen water (Mac Farlane *et al.*, 1974).

4.b Determination of Rate of Passage

A practical method to determine the rate of passage of digesta is to use an appropriate marker introduced with the feed offered and follow its appearance in the faeces. The resulting marker concentrations are plotted against time of collection, and the appropriate curve derived can be mathematically analyzed to determine the outflow rate constants. A suitable external marker is the PA6 (polyamide granules) (Becker *et al.*, 1992).

EXERCISE METABOLISM

Draft animals have been an integral part of agricultural development throughout Indonesia for centuries but mechanisation is now being introduced in some areas as a consequence of the scarcity of human labour and animal power at peak periods of land preparation. There is a great demand for additional power which could be met by providing more draft animals, especially in areas where tractors cannot operate and in transmigration areas. The large number of animals used for draft power and their importance to agricultural development in Indonesia makes it imperative that they are used efficiently by providing adequate diet. To provide the required amount of energy for optimum performance and production, it is necessary to know the energy expenditure of draft animals under the conditions in which they typically work. The actual expenditure is proportional to the type of work and work load. To estimate requirements, expenditure must be measured over a range of workloads and the predicted requirement related to work output. Expenditure for any given workload will vary with breed and will be influenced by the prevailing environmental conditions. There are presently no reliable figures from which to predict the energy requirements of draft animals in Indonesia. We used the following techniques on female swamp buffaloes in addition to balance trials.

Appropriate Methodologies on Working Animals.

1. Relation Between HR and EE of (Growing) Working Female Swamp Buffaloes

Using Polar Sport Tester, continuous long term (up to 36 hours) HR monitoring has been used successfully on swamp buffaloes in Indonesia (Mahardika *et al.*, 1995). Simultaneously calculation of energy expenditure was carried out by the factorial method of measuring work output on exercising animals. Increase in energy expenditure was related to increase in HR following the equation: $EE = 17.22 + 0.23HR$ ($r = 0.95$). Results also reveal that it is not advisable to impose a work load exceeding 15 % of liveweight on female swamp buffaloes.

The heart rate monitoring is sturdy and relatively inexpensive and our experience has shown that the necessary equipment can be used reliably in field studies. But, although the method shows great promise, it has yet to be validated in the field. For this purpose, an appropriate method would be measuring EE as the difference between ME and RE. The *in vivo* body density method was used to estimate body composition and RE.

2. Body Density (Water Displacement) Method

The swamp buffalo proved to be a suitable animal for the water displacement method. It attempted to estimate longterm EE of working buffaloes from energy balance within a fortnight experimental period and benefiting from the relationship $EE = ME - RE$. Measuring *in vivo* body composition with this body density technique showed that the fat content of buffaloes ranged from 16.8 to 18.7 %, and the protein ranged from 17.4 to 18.7 %. Treatment with 3-hours work for 14 days did not have a significant effect on body composition, although there was a tendency that the fat content decreased in the working buffaloes. Non-working buffaloes had fat retention of 0.07 kg/day, whereas working buffaloes showed negative retentions (Table 2) (Mahardika *et al.*, 1997).

Similar results also happened to protein retention. However, negative retention of protein only occurred in buffalo with 3-hours work/day. The decreased content of fat and protein resulted from the use of both substances as energy sources for work. Fat degradation would occur earlier than protein's. Both decrements of body constituents are exponential in nature. For the calculation of HP, RE should be corrected by dividing it by the efficiency coefficient for the formation of body tissue. This alternative method of HP measurement was used for the validation of the heart rate monitoring technique and will be routinely applied with ruminants in our laboratory. Equations for swamp buffaloes relating components of lean (water, bone, protein and

meat) and **fat** was made by **Mahardika et. al.** (1997) who used the data found in **Natasasmita's** PhD **dissertation** (Univ. of Melbourne, 1978). Although the method is rather **laborious**, facilities are easy to build and is not **too** costly. The method is attractive **to** be developed for other ruminant species.

Table 2 . Retentions of Fat, Protein and Energy for Working Buffaloes Subjected to Different **Working** Duration

Variables	Treatment			
	No work	1-hour work	2-hour work	3-hour work
Fat Retention (kg/day)	0.07 ^a	-0.04 ^b	-0.08 ^{bc}	-0.09 ^c
Protein Retention (kg/day)	0.11 ^a	0.08 ^a	0.01 ^b	-0.10 ^c
Energy Retention (MJ/day)	4.96 ^a	0.03 ^b	-2.75 ^c	-4.92 ^c

Note: The values with different letters on the same line are significantly different ($P < 0.05$) (Source: **Mahardika et al.**, 1997)

PREGNANT AND LACTATING RUMINANTS

The indigenous small ruminant breeds of Indonesia are adapted to the humid tropical environment. They show independence of **photoperiodicity** for **breeding** and are generally believed to be prolific. In the **free** living state, they feed on grasses or poor roughage without being given any feed supplement. ME supply with such diet is insufficient to support energy retention and ADG, the more for pregnant animals. The **conceptus** depends largely on glucose for its energy supply and demands high maternal glucose production rates (Sastradipradja *et al.*, 1994). The rapid increase in foetal growth in late pregnancy imposes a progressive limitation on the use of poor quality roughage as the sole feed. Concentrate supplementation improves ME and protein supplies (Katipana and Sastradipradja, 1994), however, such feeding needs long training for the animal's acceptance.

ME use for lactation is considered in general more efficient than for fattening and tissue gain. Total amounts of specific nutrients available within a lactating animal that are utilized for milk production, and other productive processes in the body, are not equal to amounts absorbed **from** the digestive tract. Therefore, a balance model of milk synthesis of the indigenous female is needed to evaluate carbon and nitrogen flows into and out of the gland, which generates sufficient

energy and **reducing power** to meet **synthetic** requirements. **Calculated uptakes** of **metabolites**, energy requirements of **the gland** for **synthesis** of **milk** components and **other related** quantitative **data** are needed to evaluate **effects** of changing nutrient availability and metabolic control mechanisms.

Our studies with **pregnant and** lactating small ruminants requires measurements of the same metabolic parameters as **required** for the growing animal, **e.g.** balance trials, whole **animal calorimetry**, glucose kinetics and activity of the **splanchnic** bed. For the studies on the lactating animal, additional **data** are needed on milk production and composition, and the **metabolic** activity of the mammary gland.

1. Milk Production and Composition

Prior to weaning, the development and **growth performance** of the **young** is dependent upon the milk supply **from** the mother animal, hence the amount of milk produced is an important parameter to be looked at for judging milk performance of the mother animal. Production can be measured after handmilking, usually **oxytocin** is injected prior to milking. Samples of milk can be analyzed for milk components, **e.g.** lactose, fat, **crude** protein and **citrate**. Direct measurement of milk yield may fail to provide an accurate estimate of milk production, especially for **non-dairy** animals where production is very limited. Therefore, an appropriate dilution method (deuterium oxide) for the determination of milk yield of such animals would be **useful**. The principle steps of this procedure includes injection of the **D₂O** into the body of the infant, collection of blood samples, separation of the water moiety, determination of the **D₂O** content, calculation of water turnover and calculation of water intake (Prawirodigdo and Sastradipradja, 1992).

2 Mammary Gland Metabolism

2.a Mammary Blood Flow

This physiological parameter is essential in basic trials for the calculation of nutrient **utilization** in lactating animals. Mammary blood flow (MBF) is estimated by a variety of techniques. For our purposes we choose an **indirect** technique employing the Fick principle and initially used total N in **milk**, arterial and venous blood for the calculations (**Astuti**, 1995). Later **measurements** applied a technique according to Cant *et al.* (1993) using **A-V** difference of **phenylalanine** and tyrosine, and their contents in milk protein. Thus,

$$\text{MBF (liters per hour)} = \{(FY_B \times .965) + FY_F\} / \{FY_{A-V}\},$$

where:

FY_B = Phe + Tyr output in milk protein (moles per hour),
 FY_F = free milk Phe + Tyr (moles per hour), and
 FY_{A-V} = Phe + Tyr A-V difference (moles per hour).

2.b *Metabolite Uptake, Oxygen Uptake*

Data on blood and plasma metabolite concentrations is needed and using the Fick principle, extraction percentages and uptake by the **mammary** gland can be **calculated**. Metabolites of interest are oxygen, glucose, acetate, betahydroxybutyrate (BHBA), lactate, glycerol, triglycerides (TG), other lipids, aminoacids. **Oxygen** uptake and **CO₂** production provide the basis for estimating substrate oxidation.

2.c *Biochemical Analysis*

Conventional chemical methods for the determination of concentration levels of metabolites in body fluids can provide adequate information to the **understanding** of **how** feed **nutrients** are converted into animal products and biochemical **status** of the animal. Such laboratory work includes determination of total **and** individual Volatile **Fatty Acid (VFA)**, **ruminal** ammonia-N, blood glucose, FFA, TG, alpha-keto butyric acid, urinary **allantoin**, blood plasma **aminoacids** and minerals.

3. *Lactating Dairy Cows*

Onfarm research with dairy cattle owned by commercial enterprises **suffers from limitations** imposed by the owners by not allowing major manipulations to be given to the animals or to the existing system. Under these circumstances one has to fully make use of the limited data collected. From such limited data, Sigit (1995) **managed** to construct material **balance** of lactating FH cows at the **Baruajak Farm, Bandung**.

With minimal disturbance to the animal, **data** were collected on body weight by **measuring** the chest circumference according to **Schoorl's** equation:

$$\text{BW (kg)} = 0.01[\text{chest circ. (cm)} + 22]^2,$$

feed intake as the difference between feed offered and **refused**, total collection of wastes, daily milk production and milk analysis, while data on digestibilities were obtained from *in vitro* estimations.

HP was calculated as follows:

N retention (**g/d**) was calculated as amount of N digested - **urinary N** - milk N

C retention (**g/d**) was calculated as digested C - milk C - **urinary N**

The value still contains the amount of C which would escape as respiratory **CO₂**. HP was calculated from **CO₂ production (Mcal/day)** = [retained C • C of meat and fat which disappeared due to loss of **body weight**] x **94/1211000** (Brody, 1945).

1 mole C = **12g**, produces **CO₂** with 94 Kcal. (1 Kcal = 4.185 KJ)

EXAMPLES OF RESULTS

Data of relevant metabolic parameters of growing female goats (av. BW 14 kg) and lactating does (av. BW 25 kg) fed ad libitum and restricted diets are presented in Table 3. Calculations **demonstrated** that energy maintenance requirement for **growing** goats was 0.47 **MJ/kg^{0.75}/d** while **ME_m** for the lactating doe was 0.71 **MJ/kg^{0.75}/d**. Portal blood flow ranged from 400 - 1000 **ml/min** which means about 30 % of cardiac output in growing goats. The data for lactating does were about 15 %. We found evidence that the gluconic capacity of the growing goat is adequate even with restricted feeding and similar conclusion could be drawn for the lactating doe. **Manik** and **Sastradipradja** (1989) found that **gluconeogenesis** in the lactating **goat** is high presumably from **ruminal propionate**. Metabolism of the **splanchnic** area in growing **goats** seemed to be more active than is the case with lactating does.

With JTT growing lambs, the effect of multiple clenbuterol injection (CB) was investigated. It was found that there **were, significant** quadratic responses (**P<.05**) to dose of CB (X) for average weights of 5 individual muscles (I-Y) and 5 cut yields (II-Y) of which the equations were respectively :

$$\text{I-Y} = 196.3292 + 8.5304X - 0.3164X^2 \text{ (P<.05; R}^2 = 0.8687; S_{Y..} = 10.4522). \\ \text{max. 253.8 g for CB 13.48 : g/kg BW}$$

$$\text{II-Y} = 0.8194 + 0.0393X - 0.0015 X^2 \text{ (P<.05; R}^2 = 0.7003; S_{Y..} = 0.0573). \\ \text{max. 1.069 kg for CB 12.70 : g/kg BW. (Saka et al., 1997).}$$

Table 3. Digestibility and Metabolism of Nutrients of Growing **Female** and Lactating Etawah Cross-Breed Goats

Parameter		ad <i>libitum</i>	medium	low	P level of significance
Energy/nutrient balances:					
GE (MJ/d)	G	8.53	6.27	4.71	
	L	15.98	14.07	11.38	
DE (MJ/d)	G	5.22'	3.71^{bc}	2.83'	≤ 0.01
	L	10.70'	10.12'	8.18^b	≤ 0.01
CP (g/d)	G	102	75	56	
	L	158	152	135	
ME (MJ/d)	G	4.47''	3.17^{bc}	2.40'	≤ 0.01
	L	9.26''	8.85'	7.10^b	≤ 0.01
HP (MJ/d)	G	3.62	3.25	3.31	NS
	L	6.28	5.46	5.21	NS
HP/GE (%)	G	43	51	70	
	L	39.3	38.8	45.8	
RE (MJ/d)	G	0.85'	-0.085^{ab}	-0.91 ^b	≤ 0.05
	L	3.16	3.09	1.68	NS
MBS (kg ^{0.75})	G	7.75	7.12	6.48	NS
	L	10.98	11.34	10.45	NS
Catprot(U/Nx6.25g/d)	G	12.68''	7.55^b	11.72''	≤ 0.01
	L	39.6	29.9	33.2	NS
MPS (g/d)	G	3.14'	2.65^{ab}	2.29^{ab}	≤ 0.05
	L	6.28'	4.50^b	4.14^b	≤ 0.05
RProt (g/d)	G	67.14'	48.50^b	31.01'	≤ 0.01
	L	59.95'	57.81^{ab}	37.35^b	≤ 0.05
Glu. Flux (mg/min)	G	21.02'	12.20^b	4.63^d	≤ 0.01
	L	29.43	24.20	14.46	NS
Cardiac min.vol.(ml/min)	G	3160	3160	2880	NS
	L	3670	2700	2940	NS
Organ metabolism:					
BF _{splanchnic bed} (ml/min)	G	1032	625	394	NS
	L	500'	371^b	223'	≤ 0.05
HP _{splanchnic} /HP (%)	G	45	24	16	
	L	6.3	6.8	3.7	
VFA absorp. (mM/min)	G	22.8''	15.4''	8.6^b	≤ 0.05
	L	16.7'	12.8^{ab}	7.7^b	≤ 0.05
Mammary BF (ml/min)	L	229'	193^b	128'	≤ 0.05

G = growing goats, source :Astuti *et al.* (1997)

L = lactating docs, source: Astuti (1995).

Experimental data on pregnant ewes fed grass as the sole feed revealed that the diet was insufficient and the **strains effects** of pregnancy would cause maternal fat may had been mobilized to get access to fat **glycerol** for endogenous glucose **production, eventhough** no signs of **acetonemia** were **observed** (Table 4). **RP**, **RE** and **ADG** were improved by concentrate supplementation (**Sastradipradja et al.**, 1991). The beneficial **effect** of **adequate** feeding has been demonstrated also on **pregnant** does (**Katipana and Sastradipradja**, 1994).

Table 4. Metabolic Responses of Pregnant Ewes and Does to Adequate Feeding

Parameter	R0			R1			R2		
	P0	P1	P2	RO	P1	P2	FO	P1	P2
EWES*:									
ME (MJ/d)	3-11	3.12	2.88	5.34	4.83	5.05			
HP (MJ/d)	4.22	5.85	6.08	3.75	6.00	5.53			
RE (MJ/d)	-1.12	-2.73	-3.20	1.58	-1.17	-0.48			
RP (g/d)	6.38	-3.83	-7.65	35.02	21.29	30.61			
DOES**									
ME (MJ/d)				3.09	5.99	6.06	5.51	11.98	12.18
HP (MJ/d)				2.92	4.36	4.75	2.65	6.51	9.48
RE (MJ/d)				0.17	1.64	1.31	2.86	5.47	2.6611
RP (g/d)				19.43	47.19	48.37	36.06	110.19	112.8

P0 = non-pregnant, P1 = single and P2 = twin pregnancy.

• for ewes: R0 = sole grass feed; R1 = grass + concentrate, Significant difference due to ration.

** for docs: R2 = low dietary level; R3 = medium dietary level.

Table 5. Metabolic Responses of Lactating FH Cows Fed a Ration Supplemented with **Hydroxymethionine** Analogue (**Sigit**, 1995)

Variable	A0	A1	A3
Urinary-N (g/d)	50.52	65.68	51.44
N retention (g/d)	55.7	41.88	37.05
C retention (g/d)	2488	2474	2280
ADG (g/d)	-554.7	-404.7	-781.3*
HP (MJ/d)	63.98	67.56	73.17
HP (MJ/kg ^{0.75})	0.790	0.797	0.852

A0, A1 and A3 were respectively 0, 0.1 and 0.2%. *supplementation with ammoniated zeolite at 3.0% level improved ADG (+83.33 g/d).

Table 5 contains data on **metabolic** parameters in **lactating** FH cows fed a diet supplemented with **hydroxy-methionine analogue** (AO, **A1** and **A3** were respectively 0, 0.1 and 0.2 %) (Sigit, 1995). Addition of **ammoniated zeolite** at 3.0 % improves ADG **+83.33 g/d**.

CONCLUSION

There are a number of appropriate **methodologies** available for conducting **onfarm** metabolic research **aiming** at gaining better **understanding** of animal production and health in the humid tropics.

ACKNOWLEDGEMENT

The author wishes to express his sincere gratitude to institutions which gave financial support or assistance or placed at our disposal facilities to make possible the development of laboratory techniques and the conduct of the research programs.

1. Indonesian Council of Sciences (**MIPI**), Ministry of National Research of RI, grant no. **62/Ket/1992**,
2. IAEA Technical Assistance 1978-79 Research contracts no. **1439/RB, 1863/RB**,
3. OECF Laboratory Equipment 1978-79,
4. Directorate General of Higher Education DEPDIKBUD Research contract no. **61/PSSR/DPPM/625/85, 99/P4M/DPPM/BDXXI/89, 021/P4M/DPPM/PHB/92, 021/P4M/DPPM/PHBI/2/1993, 61/P4M/DPPM/94/PHB I/3/1994, 004/P4M/DPPM/95/PHB I/4/95, 08/P2IPT/DPPM/96/PHB V/L/V/1996**,
5. Centre for Animal Research and Development AARD Indonesia,
6. Inter University Center for Life Sciences IPB.

REFERENCES

- Arta Putra, I G. , D. **Sastradipradja**, D.A. Astuti, B. Kiranadi and L.A. **Sofyan**. 1997. Nutrient Utilization of Growing Female Etawah Crossbreed Goats Fed Diets Containing Different **Protein/** Energy Ratios. *In: Energy **Metabolism** of **Farm** Animals, Proc. **14th Symp.**, Belfast UK 14-20 September 1997.*
- Astuti**, D.A., D. **Sastradipradja**, T. **Sutardi** and B. **Haryanto**. 1997. Energy and Protein Requirements of Growing Female Etawah Crossbreed Goats **Determined** From Metabolic Parameters. *In: Energy Metabolism of **Farm** Animals, Proc. **14th Symp.**, Belfast UK 14-20 **September 1997.***

- Astuti, D.A.** 1995. **Evaluasi Pemanfaatan Nutrien Berdasarkan Curahan Melalui Sistem Vena Porta dan Organ Terkait pada Kambing Tumbuh dan Laktasi.** Disertasi Doktor. Institut Pertanian Bogor.
- Becker, K., F. Mahler, E. Schlecht and A. Susenbeth. 1992. **Plant/Animal Interaction on Savanna Shrubland in Sahelian Mali.** Proceedings 5th Asia and Oceania Congr. Nucl. Med. and Biology, vol. 2, PKBNI Jakarta. **p413-434.**
- Blaxter, K.** 1989. Energy Metabolism in Farm Animals: Past, Present and Future. *In:* Y. van der Honing and **W.H. Close (eds).** Energy **Metabolism** of Farm Animals, EAAP Publ. No. 43, Centre for **Agric.** Publ. And Documentation, Pudoc Wageningen, Netherlands. **p379-384.**
- Brody, S.** 1945. Bioenergetics and Growth. **Reinhold Publ. Corp.,** New York, USA.
- Corbett, J.L., D.J. **Farrell, R.A. Leng, G.L. McClymont** and B.A. Young. 1971. **Determination of the Energy Expenditure of Penned and Grazing Sheep from Estimates of Carbondioxide Entry Rate.** *Brit. J. Nutr.,* **26:277-291.**
- Cant, **J.P, E.J. DePeters** and R.L. **Baldwin.** 1993. **Mammary Aminoacid Utilization in Dairy Cows Fed Fat and its Relationship to Milk Protein Depression.** *J. Dairy Sci.,* **76:762-774.**
- IAEA. 1985. Laboratory Training Manual on the Use of Nuclear Techniques in Animal Nutrition, **Vienna. p191-192.**
- Katipana, N.G.F.** and D. Sastradipradja. 1994. Energy and Protein Requirements of **Etawah Cross Goats** During Late Pregnancy. *In:* J.F. Aguilera (Comp.). Energy Metabolism of **Farm Animal?**, EAAP Publ. No. 76, CSIC Publ. **Serv.,** Madrid, Spain. **P.63-66.**
- Katz, M.L.** and E.N. **Bergman.** 1969. Simultaneous Measurements of Hepatic and Portal Venous Blood Flow in the Sheep and Dog. *Am. J. Physiol.,* **216:946-952.**
- Kleiber, M. 1962. The Fire of Life. An Introduction to **Animal Energetics.** John Wiley and **Sons, Inc.,** New York- London. **p44-59.**
- Macfarlane, W.V., B. Howard and B.F. Good. 1974. Tracer in Field Measurements of Water, Milk and Thyroxine Metabolism of Tropical Ruminants. *In:* Tracer Techniques in Tropical Animal Production. **IAEA, Vienna.** p. 1-23.
- Mahardika, I G., D. Sastradipradja, I K. Sumadi, T. Sutardi** and **Kartiarso.** 1995. Energy Expenditure of Water Buffaloes for Draft Power in the Humid Tropics. *In:* K. **Becker, P. Lawrence** and B. **Ørskov** (Ed). Sustainable Small-Scale Ruminant Production in Semi-Arid and Sub-Humid Tropical Areas. *Inst. An. Prod. Tropics and Subtropics, University of Hohenheim, Stuttgart, Germany.* **p127-130.**
- Mahardika, I G., D. Sastradipradja** and I K. Sumadi. 1997. Daily Energy Expenditure and Protein Requirement of Working Female Swamp Buffaloes Estimated **from** Energy Balance and

Body Composition. In: Energy Metabolism of Farm Animals, Proc. 14th Symp., Belfast UK. 14-20 September.

- Manik, I G. and D. Sastradipradja. 1989. Effect of Heat-Moisture Treated Cassava in Urea Containing Feed Supplement on Carbohydrate, Protein and Energy Metabolism of Goats. In: Y. van der Honing (Comp.). Energy Metabolism of Farm Animals, EAAP Publ. No. 43, Pudoc-Wageningen, Netherlands. **p73-76.**
- Menke, K.H. and H. Steingass. 1988. **Estmation** of the Energetic Feed Value Obtained **from** Chemical Analysis and in vitro gas production. *Anim. Res. and Dev.*, **28:7-55.**
- Natasasmita**, A. 1978. Body Composition of Swamp Buffalo. A Study of Developmental Growth and of Sex Differences. **PhD** dissertation, University of Melbourne, Australia.
- Panaretto, B.A. and A.R. Till. 1963. Body Composition In Vivo. **II.** The Composition of Mature Goats and its Relationship to the **Antipyrine**, Tritiated Water and N-acetyl-4-aminoantipyrine Spaces *Austr. J. Agric. Res.*, **14:926-943.**
- Panaretto, B.A. 1963. Body composition in **vivo**. **III.** The Composition of Living Ruminants and its Relation to the Tritiated Water Spaces. *Austr. J. Agric. Res.*, **14:944-952.**
- Prawirodigdo, S. and D. Sastradipradja. 1992. Application of the Deuterium Oxide Dilution Technique to Estimate Milk Yield of Sows. Proceedings 5 Asia and Oceania Congr. Nucl. Med. And Biology, vol. 2, PKBNI Jakarta. **p435-449.**
- Rule, D.C., R.N. Arnold, E.J. Hentges and D.C. **Beitz**. 1986. Evaluation of Urea Dilution as a Technique for Estimating Body Composition of Beef Steers In Vivo: Validation of Published Equations and Comparison with Chemical Composition. *J. Anim. Sci.*, **63:1935-1948.**
- Saka, I K., D. Sastradipradja, and B. **Kiranadi**. 1997. Nutrient Metabolism and **Carcase** Characteristics of Male Thin Tail Sheep Receiving Clenbuterol. In: Energy Metabolism of Farm Animals, Proc. 14th Symp., Belfast UK. 14-20 September.
- Sastradipradja, D. 1992. Tracer Techniques for Measuring Energy Expenditure and Related Metabolic Parameters. Proceedings 5 Asia and Oceania Congr. Nucl. Med. And Biology, vol. 2, PKBNI Jakarta. **p451-457.**
- Sastradipradja, D., N.G.F. Katipana, D.R Astuti, D.R. Ekastuti and H. **Maheshwari**. 1991. The Effect of Concentrate Supplementation on Glucose, Energy and Protein Metabolism of Single and Twin Pregnant Ewes. In: C. **Wenk** and M. Boessinger (Ed.). Energy Metabolism of Farm Animals, EAAP Publ. No. 58, **Inst. Nutztierwiss.** Gruppe Ernaehrung, **ETH-Zentrum**, Zurich, Switzerland. P. 276-279.
- Sastradipradja, D., **D.R Astuti**, H. Permadi and **P. Paridjo**. 1997 Portal Blood Flow of Female Growing and Lactating Goats on Different Food Intakes. *Media Veteriner*, **4(2): 23-33**

Sigit, N.A. 1995. Penggunaan Zeolit Beramonium dan Analog Hidroksi Metionin dalam Ransum Sapi Perah Laktasi. Disertasi Doktor. Institut Pertanian Bogor.

Suwandyastuti, S.N.O., **1986. Peningkatan Mutu Jerami Padi Ditinjau dari Neraca Mineral Esensial pada Sapi Perah. Disertasi Doktor. Institut Pertanian Bogor.**

Whitelaw, F.G., J. **Hyldgaard-Jensen**, R.S. Seid and M.G. Kay. **1970. Volatile Fatty Acid Production in the Rumen of Cattle Given an all Concentrate Diet. Brit. J. Nutr., 24:179-195.**